(FILE 'HOME' ENTERED AT 16:31:32 ON 04 DEC 2001)

FILE 'BIOSIS, EMBASE, CAPLUS, MEDLINE' ENTERED AT 16:42:48 ON 04 DEC 2001

| | | | |
|---|---|---|---|
| L1 | 2957 | S | (INTERNAL RIBOSOME ENTRY SITE SEQUENCE?) OR IRES |
| L2 | 50 | S | PITSLRE PROTEIN KINASE? |
| L3 | 253 | S | CELL CYCLE DEPENDENCY |
| L4 | 5 | S | L1 AND L2 |
| L5 | 3002 | S | L1 OR L2 |
| L6 | 0 | S | L5 AND L3 |
| L7 | 2 | DUP REM | L4 (3 DUPLICATES REMOVED) |

d l7 1-2 ti abs ibib

L7 ANSWER 1 OF 2 CAPLUS COPYRIGHT 2001 ACS

TI Cell cycle-regulated internal ribosome entry site of **PITSLRE protein kinase** gene, chimeric gene vectors for cell cycle-dependent translation initiation, and use in gene therapy

AB An internal ribosome entry site (**IRES**) of **PITSLRE protein kinase** gene, which is cell cycle-regulated, is disclosed. A method for cap-independent translation, or cell cycle-dependent translational initiation, using chimeric gene vectors contg. the **IRES**, and use in gene therapy, are also claimed. The current invention relates to two isoforms, p110 and p58 of **PITSLRE protein kinase**, which can be translated from the same p110 (.alpha.2-2) mRNA by an internal ribosome entry process. This means that p110 and p58, two proteins with putative different functions, are translated from a single mRNA species by using two AUGs within the same reading frame. These two proteins share the 439 C-terminal amino acids that contain the kinase domain. The **IRES** in the polycistronic p110 mRNA is the first **IRES** completely localized in the coding region of a cellular mRNA. Moreover, it was unexpectedly found that the **IRES** element is cell cycle regulated. Translation of p58 occurs in the G2/M stage of the cell cycle.

ACCESSION NUMBER: 2000:535275 CAPLUS
DOCUMENT NUMBER: 133:130823
TITLE: Cell cycle-regulated internal ribosome entry site of **PITSLRE protein kinase** gene, chimeric gene vectors for cell cycle-dependent translation initiation, and use in gene therapy
INVENTOR(S): Cornelis, Sigrid; Beyaert, Rudi
PATENT ASSIGNEE(S): Vlaams Interuniversitair Instituut voor Biotechnologie VZW, Belg.
SOURCE: PCT Int. Appl., 57 pp.
CODEN: PIXXD2
DOCUMENT TYPE: Patent
LANGUAGE: English
FAMILY ACC. NUM. COUNT: 1
PATENT INFORMATION:

| PATENT NO. | KIND | DATE | APPLICATION NO. | DATE |
|---|---|---|---|---|
| WO 2000044896 | A1 | 20000803 | WO 2000-EP643 | 20000126 |
| W: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM | | | | |
| RW: GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG | | | | |
| EP 1147188 | A1 | 20011024 | EP 2000-906237 | 20000126 |
| R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO | | | | |

PRIORITY APPLN. INFO.: EP 1999-200216 A 19990126
WO 2000-EP643 W 20000126

REFERENCE COUNT: 9
REFERENCE(S):
(2) Cornelis, S; MOLECULAR CELL 2000, V5(4), P597 CAPLUS
(4) Gururajan, R; GENOME RESEARCH 1998, V8, P929 CAPLUS
(5) Havenga, M; MOLECULAR IMMUNOLOGY V222(2), P319 CAPLUS

(6) Hengst, L; SCIENCE 1996, V271, P1861 CAPLUS
(8) Qbi Enterprises Ltd; WO 9821321 A 1998 CAPLUS
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L7 ANSWER 2 OF 2 BIOSIS COPYRIGHT 2001 BIOSIS DUPLICATE 1
TI Identification and characterization of a novel cell cycle-regulated internal ribosome entry site.
AB **PITSLRE protein kinases** are related to the large family of cyclin-dependent kinases. They have been proposed to act as tumor suppressor genes and have been shown to play a role in cell cycle progression. We report that two **PITSLRE protein kinase** isoforms, namely p110PITSLRE and p58PITSLRE, are translated from a single transcript by initiation at alternative in-frame AUG codons. p110PITSLRE is produced by classical cap-dependent translation, whereas p58PITSLRE results from internal initiation of translation controlled by an internal ribosome entry site (**IRES**) with unique properties. The **IRES** element is localized to the mRNA coding region, and its activity is cell cycle regulated, which permits translation of p58PITSLRE in G2/M.

ACCESSION NUMBER: 2000:266847 BIOSIS
DOCUMENT NUMBER: PREV200000266847
TITLE: Identification and characterization of a novel cell cycle-regulated internal ribosome entry site.
AUTHOR(S): Cornelis, Sigrid (1); Bruynooghe, Yanik; Denecker, Geertrui; Van Huffel, Sofie; Tinton, Sandrine; Beyaert, Rudi
CORPORATE SOURCE: (1) Department of Molecular Biology, Flanders Interuniversity Institute for Biotechnology, University of Gent, K. L. Ledeganckstraat 35, B-9000, Gent Belgium
SOURCE: Molecular Cell, (April, 2000) Vol. 5, No. 4, pp. 597-605. print..
ISSN: 1097-2765.
DOCUMENT TYPE: Article
LANGUAGE: English
SUMMARY LANGUAGE: English

=>